

A COUNTEREXAMPLE TO SCOTTISH BOOK PROBLEM 155

YOSHITO ISHIKI

ABSTRACT. We construct real Banach spaces X and Y and a bijection $U: X \rightarrow Y$ which preserves all pairwise distances on every closed ball of radius $1/4$ but is not a global isometry. The proof starts from an explicit injective map that preserves every distance at most $1/2$ and shortens one longer distance. We then enlarge the source and target through a transfinite recursion that inserts every missing target point while preserving both injectivity and this distance threshold.

1. INTRODUCTION

Scottish Book Problem 155 asks whether local preservation of distances forces global preservation for a bijection between real Banach spaces [9, Problem 155]. We use the closed ball convention from the problem. Thus, for a real Banach space X , we write $B_X(x, r) = \{z \in X \mid \|z - x\|_X \leq r\}$. The hypothesis says that for every $x \in X$ there is $r_x > 0$ such that a bijection $U: X \rightarrow Y$ preserves all pairwise distances on $B_X(x, r_x)$. The question is whether U must preserve every distance in X .

Mori proved that the answer is positive when the source space is separable, even when injectivity is replaced by surjectivity [12, Theorem 3]. Basso obtained a positive answer under the stronger metric notion of a local isometry [2, Corollary 1.3]. In that notion the local restriction must also map onto an open subset of the target. In particular, the result of Basso applies to the original hypothesis when the inverse bijection is continuous. For broader historical context on Scottish Book Problem 155, see [3, Problem 155].

The unrestricted statement fails.

Theorem 1.1. *There are real Banach spaces X and Y and a bijection $U: X \rightarrow Y$ such that*

$$\|U(p) - U(q)\|_Y = \|p - q\|_X$$

for every $x \in X$ and every $p, q \in B_X(x, 1/4)$, while U is not a global isometry.

The construction begins with an explicit piecewise linear map $S: \mathbb{R} \rightarrow \ell_1^2$. It is injective and preserves every distance at most $1/2$ but contracts one

2020 *Mathematics Subject Classification.* Primary 46B20, Secondary 54E40.

Key words and phrases. Banach space, local isometry, Lipschitz-free space, Scottish Book.

distant pair. The main difficulty is to make its image equal to an entire Banach space without losing injectivity or the fixed short distance scale.

The successor mechanism is Lemma 3.3. Starting from an injective map $V: M \rightarrow N$ and a missing point $y \in N \setminus V(M)$, it enlarges the source and target Banach spaces to $E = M \oplus_1 \mathbb{R}$ and F , and constructs a map $V^+: E \rightarrow F$. The new map puts y into its image and retains every distance up to the prescribed scale. Its injectivity coordinate is obtained from the Kuratowski embedding of the metric quotient formed by collapsing the old target N . The lemma also constructs a 1-Lipschitz linear retraction $R: F \rightarrow N$ onto the old target. The maps satisfy

$$(R \circ V^+)(m, s) = V(m)$$

whenever $|s| \leq L$. This coordinate recovery identity prevents distinct points from acquiring the same image when increasing unions are completed at limit ordinals.

After the successor and limit mechanisms are assembled, a recursion of regular cardinal length schedules every target point for entry into the image. The final unions are already complete, and the resulting map is bijective. The original contracted pair survives through linear isometric inclusions.

Organization. In Section 2, we collect the fixed short scale estimate and the facts about Lipschitz-free spaces used in the sequel. In Section 3, we establish the protected one point extension which forms the successor mechanism. In Section 4, we show how coherent 1-Lipschitz retractions with this coordinate recovery property preserve injectivity after completion at limit ordinals. In Section 5, we construct the bent map and carry out the regular cardinal recursion to prove Theorem 1.1 and formulate Problem 5.1.

Conventions and notation. All Banach spaces in this paper are real. An embedding between Banach spaces is linear and isometric.

Use of AI. An earlier version of this work was publicly posted as an *AI-Assisted Research Report* [7]. The present paper is a revised version of that report. OpenAI Codex was used in the preparation of this manuscript for language editing, L^AT_EX typesetting assistance, literature searches, and exploration of proof constructions. In particular, Codex suggested the counterexample construction and assisted in developing the protected one point extension and the coherent transfinite recursion. The author verified the arguments and takes full responsibility for the mathematical content and the final version of the manuscript. The author also understands all proofs and constructions presented in this paper.

Acknowledgements. The author was supported by JSPS KAKENHI Grant Number JP24KJ0182.

2. METRIC AND LINEAR TOOLS

We first pass from preservation at one fixed short scale to a global 1-Lipschitz estimate. We then recall the Lipschitz-free construction which turns the metric maps used later into linear maps between Banach spaces.

2.1. Maps which preserve a fixed short scale. A map $T: X \rightarrow Y$ between metric spaces is 1-Lipschitz if

$$d_Y(T(x_0), T(x_1)) \leq d_X(x_0, x_1)$$

for all $x_0, x_1 \in X$. For a linear map between Banach spaces, this is equivalent to having operator norm at most one.

Let M and N be Banach spaces, let $r > 0$, and let $V: M \rightarrow N$ satisfy

$$\|V(m_0) - V(m_1)\|_N = \|m_0 - m_1\|_M \quad \text{whenever } \|m_0 - m_1\|_M \leq r. \quad (1)$$

Fix $m_0, m_1 \in M$. Put $d = \|m_0 - m_1\|_M$, choose a positive integer n such that $d/n \leq r$, and set

$$m_k = m_0 + \frac{k}{n}(m_1 - m_0)$$

for $0 \leq k \leq n$. Each consecutive pair has distance d/n , so (1) applies to every pair m_{k-1}, m_k . The triangle inequality in N therefore yields

$$\begin{aligned} \|V(m_0) - V(m_1)\|_N &\leq \sum_{k=1}^n \|V(m_k) - V(m_{k-1})\|_N \\ &= \sum_{k=1}^n \|m_k - m_{k-1}\|_M \\ &= \|m_0 - m_1\|_M \end{aligned} \quad (2)$$

for all $m_0, m_1 \in M$. Thus every map satisfying (1) is globally 1-Lipschitz.

The successor construction uses both the sum norm and the maximum norm. We write $M \oplus_1 E$ and $M \oplus_\infty E$ for direct sums with the sum norm and the maximum norm, respectively.

We shall use the following norming form of the Hahn–Banach theorem [4, Theorem 4.4.2]. For every real Banach space X and every $x \in X$, there is $\phi \in X^*$ such that

$$\|\phi\|_{X^*} \leq 1 \quad \text{and} \quad \phi(x) = \|x\|_X.$$

A functional with these properties is called a norming functional for x .

2.2. Lipschitz-free spaces. We use Lipschitz-free spaces to build a Banach space from the glued metric space constructed in Section 3 and to turn a 1-Lipschitz retraction into a linear map. Let $(P, d, 0)$ be a pointed metric space. Let $\text{Lip}_0(P)$ be the Banach space of real Lipschitz functions $f: P \rightarrow \mathbb{R}$ which satisfy $f(0) = 0$, with norm $\|f\|_{\text{Lip}_0(P)} = \text{Lip}(f)$. For $p \in P$, let $\delta_p \in \text{Lip}_0(P)^*$ be the evaluation functional defined by

$$\delta_p(f) = f(p).$$

The Lipschitz-free Banach space over P is

$$\mathcal{F}(P) = \overline{\text{span}\{\delta_p \mid p \in P\}} \subset \text{Lip}_0(P)^*.$$

Equivalently, $\mathcal{F}(P)$ is the completion of the finitely supported signed measures on P with total mass zero under the Kantorovich–Rubinstein norm. For $\mu = \mu^+ - \mu^-$, this norm is the Wasserstein 1 distance between the equal mass measures μ^+ and μ^- .

Denote the canonical map by

$$\delta_P: P \rightarrow \mathcal{F}(P), \quad \delta_P(p) = \delta_p,$$

where $\delta_0 = 0$ and

$$\|\delta_p - \delta_q\|_{\mathcal{F}(P)} = d(p, q). \quad (3)$$

The dual space $\mathcal{F}(P)^*$ is canonically identified with $\text{Lip}_0(P)$. Every pointed 1-Lipschitz map $T: P \rightarrow X$ to a Banach space has a unique 1-Lipschitz linear map $\hat{T}: \mathcal{F}(P) \rightarrow X$ such that $\hat{T}(\delta_p) = T(p)$ for every $p \in P$. Equivalently, $\hat{T} \circ \delta_P = T$. Arens and Eells construct the molecule space for an arbitrary metric space and prove that its canonical map is isometric [1, Section 2(C) and Section 3]. For a Banach space regarded as a pointed metric space, Godefroy and Kalton record the canonical predual, the canonical isometry, and the linearization property [5, Definition 1.1, Proposition 2.1, and Lemma 2.5]. Michael later obtained a short proof of the isometric embedding in (3) using evaluation functionals and distance functions [11, Theorem].

We also use the real McShane extension theorem. A real Lipschitz function on a subset of a metric space extends to the whole space with the same Lipschitz constant [10, Theorem 1].

3. A PROTECTED ONE POINT EXTENSION

We separate the successor mechanism into its metric, linear, and recovery components. We first name the common interface used by the three components. Fix $r > 0$. A triple

$$\mathcal{A} = (M_{\mathcal{A}}, N_{\mathcal{A}}, V_{\mathcal{A}})$$

consists of real Banach spaces $M_{\mathcal{A}}$ and $N_{\mathcal{A}}$ and a map

$$V_{\mathcal{A}}: M_{\mathcal{A}} \rightarrow N_{\mathcal{A}}.$$

We call $\mathcal{A}$ a uniformly locally isometric triple at scale r if $V_{\mathcal{A}}$ is injective and

$$\|V_{\mathcal{A}}(m_0) - V_{\mathcal{A}}(m_1)\|_{N_{\mathcal{A}}} = \|m_0 - m_1\|_{M_{\mathcal{A}}} \quad (4)$$

whenever

$$\|m_0 - m_1\|_{M_{\mathcal{A}}} \leq r.$$

Let $L > 0$, let $\mathcal{A}$ be a uniformly locally isometric triple at scale r , and let

$$y \in N_{\mathcal{A}} \setminus V_{\mathcal{A}}(M_{\mathcal{A}}).$$

For another triple

$$\mathcal{B} = (M_{\mathcal{B}}, N_{\mathcal{B}}, V_{\mathcal{B}}),$$

consider maps

$$\iota_{\mathcal{A}}^{\mathcal{B}}: M_{\mathcal{A}} \rightarrow M_{\mathcal{B}}, \quad j_{\mathcal{A}}^{\mathcal{B}}: N_{\mathcal{A}} \rightarrow N_{\mathcal{B}},$$

and

$$R_{\mathcal{A}}^{\mathcal{B}}: N_{\mathcal{B}} \rightarrow N_{\mathcal{A}}.$$

We say that $\mathcal{B}$, together with these maps, is an L -protected extension of $\mathcal{A}$ at y if the first two maps are linear isometric embeddings, the third map is linear and 1-Lipschitz, and the following conditions hold.

(E1) The triple $\mathcal{B}$ is uniformly locally isometric at scale r .

(E2) We have

$$M_{\mathcal{B}} = M_{\mathcal{A}} \oplus_1 \mathbb{R}$$

and

$$\iota_{\mathcal{A}}^{\mathcal{B}}(m) = (m, 0)$$

for $m \in M_{\mathcal{A}}$.

(E3) The extension square commutes, which means that

$$V_{\mathcal{B}} \circ \iota_{\mathcal{A}}^{\mathcal{B}} = j_{\mathcal{A}}^{\mathcal{B}} \circ V_{\mathcal{A}}.$$

(E4) The embedded point

$$j_{\mathcal{A}}^{\mathcal{B}}(y)$$

belongs to

$$V_{\mathcal{B}}(M_{\mathcal{B}}).$$

(E5) The map $R_{\mathcal{A}}^{\mathcal{B}}$ is a retraction onto the old target in the sense that

$$R_{\mathcal{A}}^{\mathcal{B}} \circ j_{\mathcal{A}}^{\mathcal{B}} = \text{id}_{N_{\mathcal{A}}}.$$

(E6) For $m \in M_{\mathcal{A}}$ and $|s| \leq L$, we have

$$(R_{\mathcal{A}}^{\mathcal{B}} \circ V_{\mathcal{B}})(m, s) = V_{\mathcal{A}}(m).$$

The next lemma constructs the metric adjunction and records the properties needed for its later linearization.

Lemma 3.1. *Let M and N be real Banach spaces, let $r > 0$, and let $V: M \rightarrow N$ be injective and satisfy (1). Fix $y \in N \setminus V(M)$ and $a \in M$. Put $C = \|V(a) - y\|_N$, choose $H > C + 4r$, and set*

$$E = M \oplus_1 \mathbb{R}, \quad \iota_E: M \rightarrow E, \quad \iota_E(m) = (m, 0).$$

Define

$$A = (M \times \{0\}) \cup \{(a, H)\}$$

and $\varphi: A \rightarrow N$ by

$$\varphi(m, 0) = V(m), \quad \varphi(a, H) = y.$$

Let $P = N \cup_{\varphi} E$ be the quotient of the disjoint union of N and E obtained by identifying every $u \in A$ with $\varphi(u)$. Let

$$q: E \rightarrow P \quad \text{and} \quad \iota_N: N \rightarrow P$$

be the canonical maps. For $Z \in \{E, N\}$ and $u, v \in Z$, write $[u, v]_Z$ for the oriented affine segment from u to v . We write $C_0 * C_1$ for concatenation of

compatible paths and $\overline{\mathcal{C}_0}$ for the reversed path. An admissible polygonal path $\mathcal{C}$ in P is a finite concatenation of images under q or ι_N of affine line segments lying in E or N . If $\mathcal{C}$ has k pieces with endpoints $u_j, v_j \in Z_j$ for $1 \leq j \leq k$, where $Z_j \in \{E, N\}$, define

$$\text{len}(\mathcal{C}) = \sum_{j=1}^k \|u_j - v_j\|_{Z_j}.$$

The path metric on P is

$$d_P(z, w) = \inf\{\text{len}(\mathcal{C}) \mid \mathcal{C} \text{ is an admissible polygonal path from } z \text{ to } w\}.$$

Then φ is injective and 1-Lipschitz, ι_N is an isometric embedding, and q is injective and 1-Lipschitz. The maps satisfy

$$q \circ \iota_E = \iota_N \circ V \quad \text{and} \quad q(a, H) = \iota_N(y).$$

Moreover, if $x_0, x_1 \in E$ and $\|x_0 - x_1\|_E \leq r$, then

$$d_P(q(x_0), q(x_1)) = \|x_0 - x_1\|_E.$$

Proof. The set A is closed in E . The set A consists of the points of E that will be identified with their images under φ in the construction below. The map φ is injective because V is injective and $y \notin V(M)$. Its restriction to $M \times \{0\}$ is 1-Lipschitz by (2). For $m \in M$, the distance between $(m, 0)$ and the remaining point (a, H) satisfies

$$\begin{aligned} \|V(m) - y\|_N &\leq \|V(m) - V(a)\|_N + C \leq \|m - a\|_M + C \\ &< \|m - a\|_M + H = \|(m, 0) - (a, H)\|_E. \end{aligned}$$

Thus φ is 1-Lipschitz on A .

The definition of d_P makes q 1-Lipschitz. Let $\iota_A: A \rightarrow E$ be the inclusion. The canonical maps satisfy $q \circ \iota_A = \iota_N \circ \varphi$, as displayed in the adjunction square

$$\begin{array}{ccc} A & \xhookrightarrow{\iota_A} & E \\ \varphi \downarrow & & \downarrow q \\ N & \xrightarrow{\iota_N} & P. \end{array} \tag{5}$$

Evaluating the commuting identity on $(m, 0)$ and (a, H) yields the two identities in the statement. For readability, we write n for $\iota_N(n)$ inside P .

Every admissible polygonal path with both endpoints in E may be shortened so that it is either one segment in E or leaves E once and returns once. To see this, consider an internal segment $[u, v]_E$ whose endpoints belong to A . The corresponding segment in N satisfies

$$\begin{aligned} \text{len}([\varphi(u), \varphi(v)]_N) &= \|\varphi(u) - \varphi(v)\|_N \\ &\leq \|u - v\|_E = \text{len}([u, v]_E), \end{aligned}$$

because φ is 1-Lipschitz. Replace every internal E segment in this way. Consecutive segments in N can then be replaced by the affine segment

joining their endpoints, whose length is no larger by the triangle inequality in N . If the original path never enters N , the triangle inequality in E replaces it by the affine segment joining its endpoints. This gives the asserted two forms without increasing length. Consequently, for $x_0, x_1 \in E$,

$$\begin{aligned} d_P(q(x_0), q(x_1)) \\ = \min \left\{ \|x_0 - x_1\|_E, \right. \\ \left. \inf_{u, v \in A} (\|x_0 - u\|_E + \|\varphi(u) - \varphi(v)\|_N + \|v - x_1\|_E) \right\}. \end{aligned} \quad (6)$$

The second term on the right hand side of (6) is the infimum of the lengths of the polygonal paths $\mathcal{C}_{u,v}$ defined by

$$\mathcal{C}_{u,v} = [x_0, u]_E * [\varphi(u), \varphi(v)]_N * [v, x_1]_E.$$

Thus

$$\text{len}(\mathcal{C}_{u,v}) = \|x_0 - u\|_E + \|\varphi(u) - \varphi(v)\|_N + \|v - x_1\|_E.$$

For $n \in N$, the corresponding formula is

$$d_P(q(x_0), n) = \inf_{u \in A} (\|x_0 - u\|_E + \|\varphi(u) - n\|_N). \quad (7)$$

The same path replacement shows that ι_N is an isometric embedding. The map q is injective. Indeed, if $d_P(q(x_0), q(x_1)) = 0$, then (6) shows that either $\|x_0 - x_1\|_E = 0$ or the second term on its right hand side is zero. In the latter case, there are $u_k, v_k \in A$ such that

$$\|x_0 - u_k\|_E \rightarrow 0, \quad \|x_1 - v_k\|_E \rightarrow 0, \quad \|\varphi(u_k) - \varphi(v_k)\|_N \rightarrow 0.$$

Closedness of A places x_0 and x_1 in A . Continuity and injectivity of φ then imply $x_0 = x_1$. Thus $x_0 = x_1$ in either case.

We next show that q preserves every distance at most r . Write

$$x_0 = (m_0, s_0), \quad x_1 = (m_1, s_1), \quad d = \|x_0 - x_1\|_E \leq r.$$

For the two points of A in the second term of (6), there are three possibilities. Put

$$t = (a, H).$$

(A1) If both points belong to $M \times \{0\}$, write them as $(u, 0)$ and $(v, 0)$. Since $\|m_0 - m_1\|_M \leq d \leq r$, define the comparison path in N by

$$\mathcal{D}_{u,v} = [V(m_0), V(u)]_N * [V(u), V(v)]_N * [V(v), V(m_1)]_N.$$

Its length satisfies

$$\begin{aligned} \|m_0 - m_1\|_M &= \|V(m_0) - V(m_1)\|_N \\ &\leq \text{len}(\mathcal{D}_{u,v}) \\ &= \|V(m_0) - V(u)\|_N + \|V(u) - V(v)\|_N + \|V(v) - V(m_1)\|_N \\ &\leq \|m_0 - u\|_M + \|V(u) - V(v)\|_N + \|v - m_1\|_M. \end{aligned}$$

Here the equality uses (1), the first inequality is the triangle inequality along $\mathcal{D}_{u,v}$, and the final inequality uses (2). The corresponding path in P is

$$\mathcal{C}_{(u,0),(v,0)} = [x_0, (u, 0)]_E * [V(u), V(v)]_N * [(v, 0), x_1]_E.$$

Adding its three segment lengths gives

$$\begin{aligned} \text{len}(\mathcal{C}_{(u,0),(v,0)}) &= \|x_0 - (u, 0)\|_E + \|V(u) - V(v)\|_N + \|(v, 0) - x_1\|_E \\ &= \|m_0 - u\|_M + |s_0| + \|V(u) - V(v)\|_N + \|v - m_1\|_M + |s_1| \\ &\geq \|m_0 - m_1\|_M + |s_0 - s_1| \\ &= \|x_0 - x_1\|_E = d. \end{aligned}$$

(A2) If both points equal t , the corresponding path is

$$\mathcal{C}_{t,t} = [x_0, t]_E * [y, y]_N * [t, x_1]_E.$$

The middle segment is degenerate. Consequently, the triangle inequality in E gives

$$\begin{aligned} \text{len}(\mathcal{C}_{t,t}) &= \|x_0 - t\|_E + \text{len}([y, y]_N) + \|t - x_1\|_E \\ &= \|x_0 - t\|_E + \|t - x_1\|_E \\ &\geq \|x_0 - x_1\|_E = d. \end{aligned}$$

(A3) If one point belongs to $M \times \{0\}$ and the other equals t , write the first point as $(u, 0)$. Suppose first that the two points occur in the order $(u, 0), t$. The corresponding path in P is

$$\mathcal{C}_{(u,0),t} = [x_0, (u, 0)]_E * [V(u), y]_N * [t, x_1]_E.$$

To estimate its two E segments, consider the comparison path in E defined by

$$\mathcal{L}_u = [(u, 0), x_0]_E * [x_0, x_1]_E * [x_1, t]_E.$$

This path joins $(u, 0)$ to t . Since

$$\|(u, 0) - t\|_E = \|u - a\|_M + H \geq H,$$

the triangle inequality along $\mathcal{L}_u$ gives

$$\begin{aligned} H &\leq \|(u, 0) - t\|_E \\ &\leq \text{len}(\mathcal{L}_u) \\ &= \|x_0 - (u, 0)\|_E + \|x_0 - x_1\|_E + \|t - x_1\|_E \\ &= \|x_0 - (u, 0)\|_E + d + \|t - x_1\|_E. \end{aligned}$$

Therefore,

$$\|x_0 - (u, 0)\|_E + \|t - x_1\|_E \geq H - d.$$

Adding the three segment lengths of $\mathcal{C}_{(u,0),t}$ and using the nonnegativity of the middle length now yields

$$\begin{aligned} \text{len}(\mathcal{C}_{(u,0),t}) &= \|x_0 - (u, 0)\|_E + \|V(u) - y\|_N + \|t - x_1\|_E \\ &\geq H - d. \end{aligned}$$

Since $H > 2r \geq 2d$, this lower bound is greater than d . When the two attachment points occur in the opposite order, the corresponding path is

$$\mathcal{C}_{t,(u,0)} = [x_0, t]_E * [y, V(u)]_N * [(u, 0), x_1]_E.$$

In this order, use the comparison path

$$\tilde{\mathcal{L}}_u = [t, x_0]_E * [x_0, x_1]_E * [x_1, (u, 0)]_E.$$

It joins t to $(u, 0)$, so the same triangle inequality gives

$$\begin{aligned} H &\leq \|t - (u, 0)\|_E \\ &\leq \text{len}(\tilde{\mathcal{L}}_u) \\ &= \|x_0 - t\|_E + d + \|(u, 0) - x_1\|_E. \end{aligned}$$

It follows that

$$\begin{aligned} \text{len}(\mathcal{C}_{t,(u,0)}) &= \|x_0 - t\|_E + \|y - V(u)\|_N + \|(u, 0) - x_1\|_E \\ &\geq H - d > d. \end{aligned}$$

The cases (A1)–(A3) exhaust the second term in (6). It follows from (6) that

$$d_P(q(x_0), q(x_1)) = \|x_0 - x_1\|_E \quad (8)$$

whenever $\|x_0 - x_1\|_E \leq r$. $\square$

The next lemma turns the metric coordinate q into a coordinate taking values in a Banach space without changing the required short distances.

Lemma 3.2. *Retain the hypotheses and notation of Lemma 3.1. Use ι_N to regard N as a subset of P , and take $0 \in N \subset P$ as the distinguished basepoint of P . Let Z be the closed linear span of $\{(n, -\delta_n) \mid n \in N\}$ in $N \oplus_1 \mathcal{F}(P)$, and set*

$$B_0 = (N \oplus_1 \mathcal{F}(P))/Z,$$

with quotient map $\Pi: N \oplus_1 \mathcal{F}(P) \rightarrow B_0$. Define

$$j_0: N \rightarrow B_0, \quad j_0(n) = \Pi(n, 0),$$

and

$$\eta: P \rightarrow B_0, \quad \eta(p) = \Pi(0, \delta_p).$$

Then j_0 is a linear isometric embedding, η is 1-Lipschitz, and

$$\eta \circ \iota_N = j_0. \quad (9)$$

Moreover, if $x_0, x_1 \in E$ and $\|x_0 - x_1\|_E \leq r$, then

$$\|\eta(q(x_0)) - \eta(q(x_1))\|_{B_0} = \|x_0 - x_1\|_E. \quad (10)$$

Proof. The dual of the space before taking the quotient is

$$(N \oplus_1 \mathcal{F}(P))^* = N^* \oplus_\infty \text{Lip}_0(P).$$

For $\ell \in N^*$ and $f \in \text{Lip}_0(P)$, define the linear functional

$$\begin{aligned} \Phi_{\ell,f}: N \oplus_1 \mathcal{F}(P) &\rightarrow \mathbb{R}, \\ \Phi_{\ell,f}(n, \mu) &= \ell(n) + \mu(f). \end{aligned} \tag{11}$$

This functional annihilates Z exactly when

$$0 = \Phi_{\ell,f}(n, -\delta_n) = \ell(n) - f(n)$$

for every $n \in N$. Thus the annihilator condition is precisely $f|_N = \ell$. Whenever this condition holds, we use the same symbol for the induced functional on B_0 . In particular,

$$\Phi_{\ell,f}(\Pi(n, \mu)) = \ell(n) + \mu(f). \tag{12}$$

The dual unit ball of B_0 therefore consists of the pairs (ℓ, f) satisfying

$$\|\ell\|_{N^*} \leq 1, \quad f \in \text{Lip}_0(P), \quad \text{Lip}(f) \leq 1, \quad f|_N = \ell.$$

For $n \in N$, the Hahn–Banach theorem provides $\ell \in N^*$ such that $\|\ell\|_{N^*} \leq 1$ and $\ell(n) = \|n\|_N$. Thus ℓ is a norming functional for n . Extend ℓ to P by the McShane theorem. If f denotes this extension, then (12) implies

$$\Phi_{\ell,f}(j_0(n)) = \ell(n) = \|n\|_N.$$

The quotient map yields the reverse estimate because

$$\begin{aligned} \|j_0(n)\|_{B_0} &= \|\Pi(n, 0)\|_{B_0} \\ &= \inf_{z \in Z} \|(n, 0) + z\|_{N \oplus_1 \mathcal{F}(P)} \\ &\leq \|(n, 0)\|_{N \oplus_1 \mathcal{F}(P)} \\ &= \|n\|_N. \end{aligned}$$

The inequality uses $0 \in Z$. Hence $j_0: N \rightarrow B_0$ is a linear isometry. The map η is 1-Lipschitz because Π is 1-Lipschitz and δ_P is an isometric embedding. The quotient relation also yields (9).

We claim that $\eta \circ q$ preserves every distance at most r . Fix $x_0 = (m_0, s_0)$ and $x_1 = (m_1, s_1)$ with

$$d = \|x_0 - x_1\|_E \leq r, \quad p_i = q(x_i), \quad a_i = d_P(p_i, N).$$

By (8), we also have $d = d_P(p_0, p_1)$.

Assume first that

$$d \leq a_0 + a_1.$$

Set

$$c_0 = \min\{d, a_0\} \quad \text{and} \quad c_1 = c_0 - d.$$

Then $0 \leq c_0 \leq a_0$ and

$$|c_1| = d - c_0 = \max\{d - a_0, 0\} \leq a_1$$

because $d \leq a_0 + a_1$. Thus $|c_i| \leq a_i$ for $i \in \{0, 1\}$, and $c_0 - c_1 = d$. Define $g: N \cup \{p_0, p_1\} \rightarrow \mathbb{R}$ by $g|_N = 0$ and $g(p_i) = c_i$. The function g is 1-Lipschitz. Indeed, for $n \in N$, we have

$$|g(p_i) - g(n)| = |c_i| \leq a_i \leq d_P(p_i, n),$$

while

$$|g(p_0) - g(p_1)| = d = d_P(p_0, p_1).$$

Retain the symbol g for its McShane extension to P . The functional $\Phi_{0,g}$ belongs to the dual unit ball and satisfies

$$\Phi_{0,g}(\eta(p_0) - \eta(p_1)) = g(p_0) - g(p_1) = d.$$

It therefore yields the required lower bound for the B_0 norm of the selected pair.

Assume now that

$$d > a_0 + a_1. \quad (13)$$

By (7), we have

$$\begin{aligned} a_i &= \text{dist}_P(p_i, N) \\ &= \inf_{n \in N} d_P(p_i, n) \\ &= \inf_{n \in N} \inf_{u \in A} (\|x_i - u\|_E + \|\varphi(u) - n\|_N) \\ &= \inf_{u \in A} \|x_i - u\|_E \\ &= \text{dist}_E(x_i, A). \end{aligned}$$

For the penultimate equality, the expression inside the two infima is at least $\|x_i - u\|_E$. Conversely, for each $u \in A$, the choice $n = \varphi(u)$ makes its second term zero. The form of A shows that this distance is attained either at a point of $M \times \{0\}$ or at (a, H) . Put

$$b_i = (m_i, 0) \quad \text{and} \quad t = (a, H).$$

Since $\text{dist}_E(x_i, M \times \{0\}) = |s_i|$ is attained at b_i , there is $u_i \in \{b_i, t\}$ such that

$$a_i = \|x_i - u_i\|_E.$$

Let

$$\mathcal{P}_i = [x_i, u_i]_E$$

be the corresponding affine segment in E . Its length is a_i .

If $u_0 = u_1 = t$, then

$$\mathcal{P}_0 * \overline{\mathcal{P}_1}$$

is an admissible polygonal path from p_0 to p_1 of length $a_0 + a_1 < d$. This contradicts $d_P(p_0, p_1) = d$.

The two minimizers cannot have different forms either. After interchanging the indices if necessary, assume that $u_0 = b_0$ and $u_1 = t$. The polygonal path in E given by

$$\mathcal{R} = [b_0, x_0]_E * [x_0, x_1]_E * [x_1, t]_E$$

joins b_0 to t and has length

$$\text{len}(\mathcal{R}) = a_0 + d + a_1 < 2d \leq 2r.$$

On the other hand,

$$\|b_0 - t\|_E = \|m_0 - a\|_M + H \geq H > 4r.$$

This contradicts the triangle inequality in E . Consequently, $u_i = b_i$ for $i \in \{0, 1\}$, and hence $a_i = |s_i|$. In particular,

$$|s_0| + |s_1| < d \leq r. \quad (14)$$

For $|s| < r$ and $n \in N$, the adjunction formula reduces to

$$d_P(q(m, s), n) = |s| + \|V(m) - n\|_N. \quad (15)$$

To see this, consider the base route

$$\mathcal{B}_{m,s,n} = [(m, s), (m, 0)]_E * [V(m), n]_N$$

and the top route

$$\mathcal{T}_{m,s,n} = [(m, s), t]_E * [y, n]_N.$$

The base route has length

$$\text{len}(\mathcal{B}_{m,s,n}) = |s| + \|V(m) - n\|_N.$$

Equation (2) shows that a base route through $(u, 0)$ has length at least

$$|s| + \|m - u\|_M + \|V(u) - n\|_N \geq \text{len}(\mathcal{B}_{m,s,n}).$$

The top route satisfies

$$\text{len}(\mathcal{T}_{m,s,n}) \geq \|m - a\|_M + H - |s| + \|y - n\|_N.$$

The triangle inequality and the definition of C also imply

$$\text{len}(\mathcal{B}_{m,s,n}) \leq |s| + \|m - a\|_M + C + \|y - n\|_N.$$

It follows that

$$\text{len}(\mathcal{T}_{m,s,n}) - \text{len}(\mathcal{B}_{m,s,n}) \geq H - C - 2|s| > 0$$

because $|s| < r$ and $H > C + 4r$. Thus no route through t is shorter than $\mathcal{B}_{m,s,n}$, which proves (15).

Since $\|m_0 - m_1\|_M \leq d \leq r$, equation (1) implies

$$\|V(m_0) - V(m_1)\|_N = \|m_0 - m_1\|_M.$$

The Hahn–Banach theorem provides $\ell \in N^*$ with $\|\ell\|_{N^*} \leq 1$ such that

$$\ell(V(m_0) - V(m_1)) = \|V(m_0) - V(m_1)\|_N = \|m_0 - m_1\|_M.$$

Put $d_s = |s_0 - s_1|$ and set

$$c_0 = \min\{d_s, |s_0|\} \quad \text{and} \quad c_1 = c_0 - d_s.$$

We have $0 \leq c_0 \leq |s_0|$ and

$$|c_1| = d_s - c_0 = \max\{d_s - |s_0|, 0\} \leq |s_1|.$$

The last inequality follows from $d_s = |s_0 - s_1| \leq |s_0| + |s_1|$. Thus $c_0 - c_1 = d_s$. On $N \cup \{p_0, p_1\}$, set

$$f|_N = \ell, \quad f(p_i) = \ell(V(m_i)) + c_i.$$

For $n \in N$, the formula (15) implies

$$\begin{aligned} |f(p_i) - f(n)| &\leq \|V(m_i) - n\|_N + |c_i| \\ &\leq \|V(m_i) - n\|_N + |s_i| \\ &= d_P(p_i, n). \end{aligned}$$

Its difference on the selected pair is

$$f(p_0) - f(p_1) = \|m_0 - m_1\|_M + |s_0 - s_1| = d.$$

Hence f is 1-Lipschitz on $N \cup \{p_0, p_1\}$. Retain the symbol f for its McShane extension to P . It is an admissible dual functional. By (12), it satisfies

$$\Phi_{\ell, f}(\eta(p_0) - \eta(p_1)) = f(p_0) - f(p_1) = d.$$

Together with the 1-Lipschitz property of $\eta \circ q$, the two cases prove (10). $\square$

The two auxiliary lemmas now supply the coordinates needed for the successor mechanism. The next lemma adds a coordinate which separates points and constructs the retraction used at limit stages.

Lemma 3.3. *Let $r, L > 0$, and let*

$$\mathcal{A} = (M_{\mathcal{A}}, N_{\mathcal{A}}, V_{\mathcal{A}})$$

be a uniformly locally isometric triple at scale r . For every

$$y \in N_{\mathcal{A}} \setminus V_{\mathcal{A}}(M_{\mathcal{A}}),$$

there exist a triple $\mathcal{B}$ and maps $\iota_{\mathcal{A}}^{\mathcal{B}}$, $j_{\mathcal{A}}^{\mathcal{B}}$, and $R_{\mathcal{A}}^{\mathcal{B}}$ which make $\mathcal{B}$ an L -protected extension of $\mathcal{A}$ at y .

Proof. For the construction, write

$$M = M_{\mathcal{A}}, \quad N = N_{\mathcal{A}}, \quad V = V_{\mathcal{A}}.$$

Fix $y \in N \setminus V(M)$ and $a \in M$. Put $C = \|V(a) - y\|_N$ and choose $H > C + L + 4r$. Since $L > 0$, this choice also satisfies the height condition in Lemma 3.1. Apply Lemma 3.1 with these choices, and retain its notation E , ι_E , A , φ , P , q , and ι_N . Apply Lemma 3.2 to the resulting adjunction and retain its notation B_0 , Π , j_0 , and η .

The copy of N is closed in P . Indeed, for $x \notin A$, we have $d_P(q(x), N) = \text{dist}(x, A) > 0$. Collapse N to one point and let

$$\pi_Q: P \rightarrow Q = P/N$$

be the quotient map. Denote the collapsed point by $o = \pi_Q(0)$. Its metric is

$$d_Q(\pi_Q(p_0), \pi_Q(p_1)) = \min\{d_P(p_0, p_1), d_P(p_0, N) + d_P(p_1, N)\}.$$

By [6, Proposition 2.6], this formula defines a metric, and π_Q is 1-Lipschitz and injective on $P \setminus N$. For $z \in Q$, write

$$d_z: Q \rightarrow \mathbb{R}, \quad d_z(u) = d_Q(z, u)$$

for $u \in Q$, and let $C_b(Q)$ be the Banach space of bounded continuous real functions on Q with the supremum norm. The map $K: Q \rightarrow C_b(Q)$ defined by $K(z) = d_z - d_o$ is an isometric embedding and satisfies $K(o) = 0$ by [8, Theorem 2.6]. Let

$$G_Q = \overline{\text{span } K(Q)} \subset C_b(Q).$$

Since $K(Q) \subset G_Q$, we henceforth regard K as a map from Q to G_Q . Define $\tilde{K}: P \rightarrow G_Q$ by $\tilde{K} = K \circ \pi_Q$. The map $\tilde{K}$ is 1-Lipschitz, vanishes on N , and separates two points unless both belong to N . Set

$$F = B_0 \oplus_\infty G_Q.$$

Define

$$\iota_F: N \rightarrow F, \quad \iota_F(n) = (j_0(n), 0),$$

and

$$J: P \rightarrow F, \quad J(p) = (\eta(p), \tilde{K}(p)).$$

The forward maps are summarized by

$$\begin{aligned} P &\xrightarrow{\pi_Q} Q \xrightarrow{K} G_Q, & \tilde{K} &= K \circ \pi_Q, \\ J &= (\eta, \tilde{K}): P \rightarrow B_0 \oplus_\infty G_Q, & V^+ &= J \circ q. \end{aligned} \tag{16}$$

The first coordinate η preserves the required short distances. The second coordinate $\tilde{K}$ separates points not already separated inside the old target. The map J is 1-Lipschitz and injective. Equation (9) and the fact that $\tilde{K}$ vanishes on N show that

$$J \circ \iota_N = \iota_F. \tag{17}$$

The first coordinate and (10) show that $J \circ q$ preserves every distance at most r . By (16), V^+ is injective.

$$\begin{array}{ccc} M & \xrightarrow{\iota_E} & E \\ V \downarrow & & \downarrow V^+ \\ N & \xrightarrow{\iota_F} & F. \end{array} \tag{18}$$

The square commutes because $q \circ \iota_E = \iota_N \circ V$, and equation (17) then implies commutativity. The distinguished point at height H also satisfies

$$V^+(a, H) = J(\iota_N(y)) = \iota_F(y).$$

Set

$$\mathcal{B} = (E, F, V^+), \quad \iota_{\mathcal{A}}^{\mathcal{B}} = \iota_E, \quad j_{\mathcal{A}}^{\mathcal{B}} = \iota_F.$$

The construction so far proves (E1)–(E4).

It remains to prove (E5) and (E6). Let $\lambda: \mathbb{R} \rightarrow [0, 1]$ vanish on $(-\infty, L]$, be affine on $[L, H]$, and equal one on $[H, \infty)$. Define

$$\rho_E: E \rightarrow N, \quad \rho_E(m, s) = V(m) + \lambda(s)(y - V(a)).$$

Since

$$\text{Lip}(\lambda) = \frac{1}{H - L} \quad \text{and} \quad \frac{C}{H - L} < 1,$$

we have

$$\begin{aligned} & \|\rho_E(m_0, s_0) - \rho_E(m_1, s_1)\|_N \\ & \leq \|V(m_0) - V(m_1)\|_N + C|\lambda(s_0) - \lambda(s_1)| \\ & \leq \|m_0 - m_1\|_M + \frac{C}{H - L}|s_0 - s_1| \\ & \leq \|m_0 - m_1\|_M + |s_0 - s_1|. \end{aligned}$$

Therefore, the map $\rho_E: E \rightarrow N$ is 1-Lipschitz. It agrees with φ on A . Together with the identity on N , it descends to a pointed 1-Lipschitz retraction $\rho_P: P \rightarrow N$. By construction,

$$\rho_P \circ q = \rho_E, \quad \rho_P \circ \iota_N = \text{id}_N. \quad (19)$$

By the universal property of $\mathcal{F}(P)$, the map ρ_P has a unique 1-Lipschitz linear extension

$$\widehat{\rho}_P: \mathcal{F}(P) \rightarrow N.$$

It satisfies

$$\widehat{\rho}_P(\delta_p) = \rho_P(p)$$

for every $p \in P$. Define

$$T: N \oplus_1 \mathcal{F}(P) \rightarrow N, \quad T(n, \mu) = n + \widehat{\rho}_P(\mu).$$

For every $n \in N$ and $\mu \in \mathcal{F}(P)$, we have

$$\|T(n, \mu)\|_N \leq \|n\|_N + \|\widehat{\rho}_P(\mu)\|_N \leq \|n\|_N + \|\mu\|_{\mathcal{F}(P)}.$$

Thus T is 1-Lipschitz. Since $\rho_P(n) = n$ for $n \in N$, we have

$$T(n, -\delta_n) = n - \widehat{\rho}_P(\delta_n) = n - \rho_P(n) = 0.$$

It follows that $Z \subset \ker T$. Consequently, T induces a 1-Lipschitz linear map

$$R_0: B_0 \rightarrow N, \quad R_0(\Pi(n, \mu)) = n + \widehat{\rho}_P(\mu).$$

For $n \in N$, we have

$$R_0(j_0(n)) = n.$$

Hence R_0 is a linear retraction onto the old target. Moreover, the defining formula yields

$$R_0 \circ \eta = \rho_P. \quad (20)$$

Define the first coordinate projection by

$$\pi_{B_0}: F \rightarrow B_0, \quad \pi_{B_0}(b, g) = b.$$

Set

$$R = R_0 \circ \pi_{B_0}: F \rightarrow N.$$

This is a 1-Lipschitz linear map, and $R \circ \iota_F = \text{id}_N$. The named maps now fit into the commutative recovery diagram

$$\begin{array}{ccccc}
 E & \xrightarrow{q} & P & \xrightarrow{J} & F \\
 \rho_E \downarrow & & \rho_P \downarrow & & \downarrow R \\
 N & \xlongequal{\quad} & N & \xlongequal{\quad} & N.
 \end{array} \tag{21}$$

The left square is (19). The right square follows from (20) because $\pi_{B_0} \circ J = \eta$. Consequently,

$$(R \circ V^+)(m, s) = \rho_E(m, s) = V(m) + \lambda(s)(y - V(a)).$$

If $|s| \leq L$, then $\lambda(s) = 0$, and therefore

$$(R \circ V^+)(m, s) = V(m).$$

Finally, set

$$R_{\mathcal{A}}^{\mathcal{B}} = R.$$

The preceding identities prove (E5) and (E6), which completes the proof. $\square$

4. COHERENT LIMIT STAGES

We record how the retractions in Lemma 3.3 control completed increasing unions. Preservation at the fixed distance scale alone does not ensure that the extension to a completed increasing union remains injective. The retractions supply the missing coordinate recovery.

Consider a transfinite chain $\{(M_\alpha, N_\alpha, V_\alpha)\}_{\alpha < \tau}$ of Banach spaces and injective maps. For $\alpha \leq \beta < \tau$, let

$$\iota_\alpha^\beta: M_\alpha \rightarrow M_\beta \quad \text{and} \quad j_\alpha^\beta: N_\alpha \rightarrow N_\beta$$

be linear isometric embeddings. These stage embeddings are required to satisfy

$$\iota_\alpha^\alpha = \text{id}_{M_\alpha}, \quad \iota_\beta^\gamma \circ \iota_\alpha^\beta = \iota_\alpha^\gamma$$

and

$$j_\alpha^\alpha = \text{id}_{N_\alpha}, \quad j_\beta^\gamma \circ j_\alpha^\beta = j_\alpha^\gamma$$

whenever $\alpha \leq \beta \leq \gamma < \tau$. We also require the square

$$\begin{array}{ccc}
 M_\alpha & \xrightarrow{\iota_\alpha^\beta} & M_\beta \\
 V_\alpha \downarrow & & \downarrow V_\beta \\
 N_\alpha & \xrightarrow{j_\alpha^\beta} & N_\beta
 \end{array} \tag{22}$$

to commute. Thus

$$V_\beta \circ \iota_\alpha^\beta = j_\alpha^\beta \circ V_\alpha. \tag{23}$$

We use the stage embeddings to identify each stage with its image in every later stage. Under these identifications, the source spaces and target spaces

are increasing chains and the stage maps extend one another. Assume that every $V_\alpha: M_\alpha \rightarrow N_\alpha$ preserves distances at most a fixed $r > 0$. Fix $L > 0$. Whenever Lemma 3.3 is applied at a successor stage $\alpha + 1$, set

$$M_{\alpha+1} = M_\alpha \oplus_1 \mathbb{R},$$

take $\iota_\alpha^{\alpha+1}$ and $j_\alpha^{\alpha+1}$ to be the embeddings ι_E and ι_F from that lemma, and let

$$R_\alpha^{\alpha+1}: N_{\alpha+1} \rightarrow N_\alpha$$

be the 1-Lipschitz linear retraction obtained there. It satisfies

$$(R_\alpha^{\alpha+1} \circ V_{\alpha+1})(m, s) = V_\alpha(m) \quad (24)$$

whenever $|s| \leq L$. At a successor stage where no extension is required, set

$$M_{\alpha+1} = M_\alpha, \quad N_{\alpha+1} = N_\alpha, \quad V_{\alpha+1} = V_\alpha,$$

and use the identity as $\iota_\alpha^{\alpha+1}$, $j_\alpha^{\alpha+1}$, and $R_\alpha^{\alpha+1}$.

Retractions to earlier stages are defined by composition at successors. They satisfy

$$R_\alpha^\beta \circ j_\alpha^\beta = \text{id}_{N_\alpha}, \quad R_\alpha^\gamma \circ j_\beta^\gamma = R_\alpha^\beta \quad (25)$$

whenever $\alpha \leq \beta \leq \gamma$.

Proposition 4.1. *Let λ be a limit stage of the construction above. Assume that every stage map below λ is injective and preserves every distance at most r . Let M_λ and N_λ be the completions of the corresponding algebraic increasing unions. Then the coherent union of the stage maps extends uniquely to an injective 1-Lipschitz map*

$$V_\lambda: M_\lambda \rightarrow N_\lambda$$

which extends every earlier stage map and preserves every distance at most r . For each $\alpha < \lambda$, the coherent retractions extend uniquely to a 1-Lipschitz linear retraction

$$R_\alpha^\lambda: N_\lambda \rightarrow N_\alpha$$

and continue to satisfy (25).

More precisely, call the transition from stage η to stage $\eta + 1$ active when it applies Lemma 3.3. For $\delta \leq \lambda$, define

$$I(\delta) = \{\eta < \delta \mid \text{the transition from } \eta \text{ to } \eta + 1 \text{ is active}\}.$$

Thus each $\eta \in I(\delta)$ indexes the new $\mathbb{R}$ coordinate introduced in that transition, while an idle transition contributes no coordinate. We have $I(\gamma) = I(\lambda) \cap \gamma$ for $\gamma < \lambda$. Under the stage embeddings, the source spaces have the canonical coordinate representations

$$M_\gamma = M_0 \oplus_1 \ell_1(I(\gamma))$$

for $\gamma \leq \lambda$. In particular,

$$M_\lambda = M_0 \oplus_1 \ell_1(I(\lambda)).$$

For $\gamma < \lambda$, let

$$P_\gamma: M_\lambda \rightarrow M_\gamma$$

be the projection onto stage γ , defined by

$$P_\gamma(m, (s_\eta)_{\eta \in I(\lambda)}) = (m, (s_\eta)_{\eta \in I(\gamma)}).$$

If

$$x = (m, (s_\eta)_{\eta \in I(\lambda)})$$

and γ is chosen so that $\eta < \gamma$ whenever $|s_\eta| > L$, then

$$(R_\gamma^\lambda \circ V_\lambda)(x) = V_\gamma(P_\gamma(x)). \quad (26)$$

Proof. Fix a limit stage λ and assume by transfinite induction that the recovery identity has been established at every earlier limit stage. For fixed $\alpha < \lambda$, the maps R_α^β agree on overlaps by (25). They therefore define a 1-Lipschitz linear map on the algebraic union by

$$R_\alpha^\lambda(j_\beta^\lambda(n)) = R_\alpha^\beta(n)$$

for $\alpha \leq \beta < \lambda$ and $n \in N_\beta$. This map extends uniquely to a 1-Lipschitz linear retraction $R_\alpha^\lambda: N_\lambda \rightarrow N_\alpha$. The extensions continue to satisfy (25).

At every transition indexed by $\eta \in I(\lambda)$, the protected extension replaces the current source M_η by $M_{\eta+1} = M_\eta \oplus_1 \mathbb{R}$, and the stage embedding satisfies

$$\iota_\eta^{\eta+1}(m) = (m, 0)$$

for m in M_η . An idle transition leaves the source unchanged. Induction over the stages below λ therefore identifies M_α with $M_0 \oplus_1 \ell_1(I(\alpha))$ inside $M_0 \oplus_1 \ell_1(I(\lambda))$ by extending later coordinates by zero. Consequently,

$$M_0 \oplus_1 c_{00}(I(\lambda)) \subseteq \bigcup_{\alpha < \lambda} M_\alpha \subseteq M_0 \oplus_1 \ell_1(I(\lambda)),$$

where $c_{00}(I(\lambda))$ denotes the finitely supported scalar families on $I(\lambda)$ with the ℓ_1 norm. The first inclusion holds because every finite subset of $I(\lambda)$ is contained in $I(\gamma)$ for some $\gamma < \lambda$. Since the completion of $c_{00}(I(\lambda))$ in this norm is $\ell_1(I(\lambda))$, the algebraic increasing union is dense in the right hand side. Therefore, the completed source has the canonical form

$$M_\lambda = M_0 \oplus_1 \ell_1(I(\lambda)). \quad (27)$$

The coherent union of the maps V_α is 1-Lipschitz by (2). It therefore extends uniquely to a 1-Lipschitz map

$$V_\lambda: M_\lambda \rightarrow N_\lambda.$$

For every $\alpha < \lambda$, the extension satisfies

$$V_\lambda \circ \iota_\alpha^\lambda = j_\alpha^\lambda \circ V_\alpha.$$

Common finite coordinate truncations show that V_λ preserves every distance at most r . For a finite set $F \subset I(\lambda)$, let

$$\tau_F: M_\lambda \rightarrow M_\lambda$$

be the coordinate projection which retains the M_0 component and the coordinates indexed by F . Its operator norm satisfies

$$\|\tau_F\|_{\text{op}} \leq 1.$$

If $x_0, x_1 \in M_\lambda$ and $\|x_0 - x_1\|_{M_\lambda} \leq r$, then $\tau_F(x_0)$ and $\tau_F(x_1)$ belong to a common earlier stage and remain at distance at most r . Their distance is therefore preserved by that stage map and hence by V_λ . As the finite sets F increase, $\tau_F(x_i)$ converges to x_i for $i \in \{0, 1\}$. Continuity now yields equality at the limit.

Fix $x \in M_\lambda$ and write

$$x = (m, (s_\alpha)_{\alpha \in I(\lambda)}) \in M_0 \oplus \ell_1(I(\lambda)).$$

Choose $\gamma < \lambda$ so that $\alpha < \gamma$ whenever $\alpha \in I(\lambda)$ and $|s_\alpha| > L$. There are only finitely many such coordinates because the tail in (27) belongs to $\ell_1(I(\lambda))$.

For a finite set $F \subset I(\lambda) \cap [\gamma, \lambda)$, let

$$\tau_{\gamma, F}: M_\lambda \rightarrow M_\lambda$$

be the coordinate projection which retains the M_0 component and the coordinates indexed by $I(\gamma) \cup F$. Put $x_F = \tau_{\gamma, F}(x)$. Choose δ with $\gamma \leq \delta < \lambda$ so that $x_F \in M_\delta$. By compatibility of the stage maps and (25), we have

$$V_\lambda(x_F) = j_\delta^\lambda(V_\delta(x_F)) \quad \text{and} \quad R_\gamma^\lambda \circ j_\delta^\lambda = R_\gamma^\delta.$$

Every coordinate of x_F added after stage γ has absolute value at most L . The successor identity (24), the identity maps at idle stages, and the induction hypothesis at earlier limit stages therefore yield the complete map calculation

$$\begin{aligned} (R_\gamma^\lambda \circ V_\lambda)(x_F) &= (R_\gamma^\lambda \circ j_\delta^\lambda \circ V_\delta)(x_F) \\ &= (R_\gamma^\delta \circ V_\delta)(x_F) \\ &= V_\gamma(P_\gamma(x)). \end{aligned} \tag{28}$$

As F increases through the finite subsets of $I(\lambda) \cap [\gamma, \lambda)$, the points x_F converge to x . Continuity therefore yields (26).

Assume that

$$V_\lambda(x_0) = V_\lambda(x_1).$$

Write

$$x_i = (m_i, (s_{i,\eta})_{\eta \in I(\lambda)})$$

for $i \in \{0, 1\}$. Choose $\beta < \lambda$ so that $\eta < \beta$ whenever $\eta \in I(\lambda)$ and

$$\max\{|s_{0,\eta}|, |s_{1,\eta}|\} > L.$$

For every γ with $\beta \leq \gamma < \lambda$, apply R_γ^λ to the equality and use (26). We obtain

$$V_\gamma(P_\gamma(x_0)) = V_\gamma(P_\gamma(x_1)).$$

Injectivity at stage γ implies

$$P_\gamma(x_0) = P_\gamma(x_1).$$

The projection onto every stage retains the M_0 component. For each $\eta \in I(\lambda)$, choose γ with $\max\{\beta, \eta + 1\} \leq \gamma < \lambda$. The equality of the two stage projections then implies $s_{0,\eta} = s_{1,\eta}$. Hence $x_0 = x_1$. $\square$

Together with Lemma 3.3, Proposition 4.1 shows by transfinite induction that all successor maps and all completed limit maps remain injective and preserve every distance at most r .

5. THE COUNTEREXAMPLE

We now assemble the counterexample. The construction has three parts. We first define the initial map, then control the size of every stage so that target points can be scheduled, and finally run the transfinite recursion.

5.1. The bent seed. Define the polygonal map $S: \mathbb{R} \rightarrow \ell_1^2$ by

$$S(t) = \begin{cases} (t, 0), & t \leq 0, \\ (0, t), & 0 \leq t \leq 1, \\ (1 - t, 1), & 1 \leq t. \end{cases} \quad (29)$$

The image consists of two parallel rays joined by a vertical segment, so S is injective. If $|s - t| \leq 1/2$, the parameter interval between s and t meets at most one corner. At either corner the two directions occupy different ℓ_1 coordinates. It follows that

$$\|S(s) - S(t)\|_{\ell_1^2} = |s - t| \quad (30)$$

whenever $|s - t| \leq 1/2$. On the other hand,

$$\|S(-1) - S(2)\|_{\ell_1^2} = 1 < 3 = |-1 - 2|. \quad (31)$$

5.2. Cardinal bounds and scheduling. The bent map supplies the required failure of global distance preservation, but its image is not the whole target. To add every missing target point later, we first choose a recursion length larger than the cardinality of every possible stage. Put

$$\mathfrak{c} = 2^{\aleph_0}, \quad \theta = 2^{\mathfrak{c}}, \quad \kappa = \theta^+.$$

The cardinal κ is regular and uncountable. We shall construct a chain through all ordinals below κ .

Every stage of the construction will have cardinality at most θ . The initial spaces have this property. At a successor stage, the metric adjunction constructed in Lemma 3.1 has cardinality at most θ . Its Lipschitz-free space has density at most θ . The relative Banach envelope B_0 from Lemma 3.2 therefore has density at most θ . The metric quotient Q has cardinality at most θ , and its Kuratowski image has closed linear span G_Q of density at most θ . Thus the direct sum used in Lemma 3.3 also has density at most θ . A Banach space of density at most θ has cardinality at most

$$\theta^{\aleph_0} = (2^{\mathfrak{c}})^{\aleph_0} = 2^{\mathfrak{c}} = \theta.$$

If $\lambda < \kappa$ is a limit ordinal, then $|\lambda| \leq \theta$. The increasing union at stage λ therefore has cardinality at most θ , and its completion has the same bound. This proves the stage bound by transfinite induction.

To ensure that every target point is eventually included in the image, we use a standard bookkeeping argument. Fix an injection

$$\sigma: \kappa \times \kappa \rightarrow \kappa \quad (32)$$

such that $\alpha < \sigma(\alpha, \xi)$ for all $\alpha, \xi < \kappa$. Such an injection can be constructed recursively after well ordering $\kappa \times \kappa$ in order type κ . At each step fewer than κ ordinals have been used, while the tail above the selected first coordinate still has cardinality κ .

5.3. Proof of the main theorem. At a successor transition which processes a missing target point, we use Lemma 3.3. An idle successor transition uses the identity maps, and a limit stage uses Proposition 4.1. Equation (23) then makes the stage maps into one well-defined final map.

Proof of Theorem 1.1. We construct the stage data by transfinite recursion on $\beta < \kappa$. When stage β is formed, the data consist of Banach spaces M_β and N_β , a map $V_\beta: M_\beta \rightarrow N_\beta$, and a surjection

$$e_\beta: \kappa \rightarrow N_\beta.$$

We retain the constants

$$r = 1/2 \quad \text{and} \quad L = 1$$

at every stage. The stage invariant is that V_β is injective and preserves every distance at most r .

At stage zero, set

$$M_0 = \mathbb{R}, \quad N_0 = \ell_1^2, \quad V_0 = S,$$

where S is defined by (29). The cardinal bound permits us to choose the surjection

$$e_0: \kappa \rightarrow N_0.$$

Assume next that stage β has been formed. Once stage δ has been formed, the pair $(\delta, \xi) \in \kappa \times \kappa$ represents the requirement that the point $e_\delta(\xi)$ eventually belong to the image of a stage map. The schedule map assigns this $\kappa \times \kappa$ -indexed family of requirements to distinct transition indices below κ . At the transition from β to $\beta + 1$, there is at most one pair (δ, ξ) with

$$\beta = \sigma(\delta, \xi)$$

because σ is injective. If there is no such pair, leave the stage unchanged. If there is such a pair, then $\delta < \beta$ by the defining property of σ . The stage embedding therefore defines the scheduled point

$$y_{\delta, \xi}^\beta := j_\delta^\beta(e_\delta(\xi)) \in N_\beta.$$

When $y_{\delta, \xi}^\beta \notin V_\beta(M_\beta)$, the stage invariant permits us to apply Lemma 3.3 to V_β and $y_{\delta, \xi}^\beta$. When $y_{\delta, \xi}^\beta$ already belongs to the image, leave the stage

unchanged. After $N_{\beta+1}$ has been formed, the cardinal bound permits us to choose

$$e_{\beta+1}: \kappa \rightarrow N_{\beta+1}.$$

At a limit ordinal $\lambda < \kappa$, form M_λ , N_λ , and V_λ as in Proposition 4.1. The cardinal bound then permits us to choose

$$e_\lambda: \kappa \rightarrow N_\lambda.$$

This completes the transfinite recursion.

The same recursive clauses prove by transfinite induction that every V_β is injective and preserves every distance at most $1/2$. At an active successor transition this follows from Lemma 3.3. At an idle transition it is immediate, and at a limit stage it follows from Proposition 4.1. The stage maps satisfy (23), and every stage embedding is a linear isometry. The cardinal calculation above verifies at each recursive clause that the required enumeration exists.

Define the final spaces by

$$X = \bigcup_{\alpha < \kappa} M_\alpha, \quad Y = \bigcup_{\alpha < \kappa} N_\alpha.$$

Define

$$U: X \rightarrow Y, \quad U(x) = V_\alpha(x) \quad \text{when } x \in M_\alpha.$$

Equation (23) shows that this definition is independent of the chosen stage α . The spaces X and Y are complete. Indeed, the members of a Cauchy sequence $\{x_n\}_{n \in \mathbb{Z}_{\geq 0}}$ belong to countably many stages. Their stage indices have supremum below κ by regularity. The entire sequence is therefore contained in one earlier Banach stage and converges there. The same argument applies to the target.

The map U is injective and preserves every distance at most $1/2$, since every pair of source points belongs to a common stage. It is also surjective. To see this, fix $y \in Y$. There is $\delta < \kappa$ with $y \in N_\delta$, and there is $\xi < \kappa$ with

$$e_\delta(\xi) = y.$$

Put $\beta = \sigma(\delta, \xi)$. Under the stage identifications, the scheduled point

$$y_{\delta, \xi}^\beta = j_\delta^\beta(y)$$

is the same element of Y as y . At the transition from β to $\beta + 1$, this point is put into the image if it is still missing. It remains in the image at every later stage.

If $p, q \in B_X(x, 1/4)$, then $\|p - q\|_X \leq 1/2$. Hence

$$\|U(p) - U(q)\|_Y = \|p - q\|_X$$

by the fixed short distance property. Finally, the points -1 and 2 from the initial source remain at distance three, while their images remain at distance one by (31). Thus U is not a global isometry. $\square$

The counterexample shows that the local closed ball hypothesis alone is insufficient. The positive results recalled in the introduction leave the following problem.

Problem 5.1. Find further structural conditions on real Banach spaces X and Y , or regularity conditions on a bijection $U: X \rightarrow Y$, which ensure that the local closed ball hypothesis of Scottish Book Problem 155 forces U to be a global isometry.

REFERENCES

- [1] R. F. Arens and J. Eells Jr. *On embedding uniform and topological spaces*. *Pacific Journal of Mathematics* 6.3 (1956), pp. 397–403. DOI: 10.2140/pjm.1956.6.397.
- [2] G. Basso. *On a problem of Mazur and Sternbach*. 2023. arXiv: 2308.08400 [math.MG].
- [3] S. Domoradzki, M. Stawiska, and M. Zarichnyi. *90+ years of the Scottish Book*. 2026. arXiv: 2603.27867 [math.HO].
- [4] H. Geiss and S. Geiss. *Measure, Probability and Functional Analysis*. Universitext. Cham: Springer, 2025. DOI: 10.1007/978-3-031-84067-8.
- [5] G. Godefroy and N. J. Kalton. *Lipschitz-free Banach spaces*. *Studia Mathematica* 159.1 (2003), pp. 121–141. DOI: 10.4064/sm159-1-6.
- [6] Y. Ishiki. *A factorization of metric spaces*. *Colloq. Math.* 174.1 (2023), pp. 101–119. DOI: 10.4064/cm9066-6-2023.
- [7] Y. Ishiki. *AI-Assisted Research Report: A counterexample to Scottish Book Problem 155*. AI-Assisted Research Report. Version 1.0. Zenodo, Aug. 30, 2026. DOI: 10.5281/zenodo.22168214. URL: <https://yoshito-ishiki-math.github.io/ai-research-reports.html> (visited on 08/30/2026).
- [8] Y. Ishiki. *An interpolation of metrics and spaces of metrics*. *Topology Appl.* 388, 109856 (2026). DOI: 10.1016/j.topol.2026.109856.
- [9] R. D. Mauldin, ed. *The Scottish Book. Mathematics from the Scottish Café, with Selected Problems from the New Scottish Book*. 2nd ed. Cham: Birkhäuser/Springer, 2015. DOI: 10.1007/978-3-319-22897-6.
- [10] E. J. McShane. *Extension of range of functions*. *Bulletin of the American Mathematical Society* 40.12 (1934), pp. 837–842. DOI: 10.1090/S0002-9904-1934-05978-0.
- [11] E. Michael. *A short proof of the Arens–Eells embedding theorem*. *Proceedings of the American Mathematical Society* 15.3 (1964), pp. 415–416. DOI: 10.1090/S0002-9939-1964-0162222-5.
- [12] M. Mori. *On the Scottish Book Problem 155 by Mazur and Sternbach*. *Comptes Rendus. Mathématique* 362 (2024), pp. 813–816. DOI: 10.5802/crmath.572.

DEPARTMENT OF MATHEMATICAL SCIENCES, TOKYO METROPOLITAN UNIVERSITY,
MINAMI-OSAWA, HACHIOJI, TOKYO 192-0397, JAPAN
Email address: `ishiki-yoshito@tmu.ac.jp`